

REFERENCES (continued)

2004. Atlanta: US Department of Health and Human Services, CDC; 2005.

10. Substance Abuse and Mental Health Services Administration. 2004 National Survey on Drug Use & Health.

11. Leigh BC, Stall R. Substance use and risky sexual behavior for exposure to HIV: issues in methodology, interpretation, and prevention. *American Psychologist* 1993;48:1035-1045.

12. The Kaiser Family Foundation. National Survey of Teens on HIV/AIDS, 2000.

13. US Census Bureau. Poverty: 1999. Census 2000 Brief. May 2003.

14. Diaz T, Chu SY, Buehler JW, et al. Socioeconomic differences among people with AIDS: results from a multistate surveillance project. *American Journal of Preventive Medicine* 1994;10:217-222.

15. Office of the Surgeon General. The Surgeon General's call to action to promote sexual health and responsible sexual behavior, July 9, 2001.

16. Clark LF, Miller KS, Nagy SS, et al. Adult identity mentoring: reducing sexual risk for African-American seventh grade students. *Journal of Adolescent Health* 2005;37:337.e1-337.e10.

17. Dittus P, Miller KS, Kotchick BA, Forehand R. Why Parents Matter!: the conceptual basis for a community-based HIV prevention program for the parents of African American youth. *Journal of Child and Family Studies* 2004;13(1):5-20.



Jackson State University
MISSISSIPPI URBAN RESEARCH CENTER

Youths, Adolescents and Young Adults Center
350 W. Woodrow Wilson Blvd. • Jackson, MS 39213
JSU Box 17309 • Jackson, MS 39217

Skills-Building Course Calendar

For more information about upcoming capacity-building assistance skills-building course offerings, please call the Youths, Adolescents and Young Adults Center of the African American Prevention Intervention Network, toll-free, at: 1-866-JSU-MURC (578-6872).



Volume 1, Issue 1

JACKSON STATE UNIVERSITY

Spring 2007



Akoma



Youths, Adolescents and Young Adults Center of the African American Prevention Intervention Network

Inspiration

i wish

i wish people were happy and secure with themselves and not feel the need to project an image of something that they are not

i wish people didn't need the approval of other people to be themselves

be who you want and what you want to be

just don't hurt others being you

i wish people didn't project an image that is not true

because when you project yourself to be more intelligent, knowledgeable and spiritual than others, ultimately your ignorance shines through brighter, louder, and clearer than the image you've projected

i wish people could just be themselves

~ Lorenzo C. Robertson

CONFERENCES

2007 HIV Prevention Leadership Summit (HPLS)

New Orleans, La. ... May 20-23, 2007

2007 United States Conference on AIDS (USCA)

Palm Springs, Calif. ... Nov. 7-10, 2007

INSIDE

2...HIV/AIDS among Youth continued

3...HIV/AIDS among Youth continued

4...Skills-Building Course Calendar

HIV/AIDS among Youth

Young people in the United States are at persistent risk for HIV infection. This risk is especially notable for youth of minority races and ethnicities. Continual HIV prevention outreach and education efforts, including programs on abstinence and on delaying the initiation of sex, are required as new generations replace the generations that benefited from earlier prevention strategies. Unless otherwise noted, this fact sheet defines youth, or young people, as persons who are 13-24 years of age.

STATISTICS

HIV/AIDS in 2004

The following are based on data from the 35 areas with long-term, confidential name-based HIV reporting.

- An estimated 4,883 young people received a diagnosis of HIV infection or AIDS, representing about 13% of the persons given a diagnosis during that year [1].

- HIV infection progressed to AIDS more slowly among young people than among all persons with a diagnosis of HIV infection. The following are the proportions of persons in whom HIV infection did not progress to AIDS within 12 months after diagnosis of HIV infection:


- 81% of persons aged 15-24
- 70% of persons aged 13-14
- 61% of all persons

- African Americans were disproportionately affected by HIV infection, accounting for 55% of all HIV infections reported among persons aged 13-24 [2].

- Young men who have sex with men (MSM), especially those of minority races or ethnicities, were at high risk for HIV infection. In the 7 cities that participated in CDC's Young Men's Survey during 1994-1998, 14% of African American MSM and 7% of Hispanic MSM aged 15-22 were infected with HIV [3].

- During 2001-2004, in the 33 states with long-term, confidential name-based HIV reporting, 62% of the 17,824 persons 13-24 years of age given a diagnoses of HIV/AIDS were males, and 38% were females.

Age of persons with HIV infection or AIDS diagnosed during 2004



Note. Based on data from 35 areas with long-term, confidential name-based HIV reporting.

- An estimated 2,174 young people received a diagnosis of AIDS (5.1% of

Continued on page 2



ADMINISTRATIVE CORE

Velesha P. Williams
Project Director

Helenrine Seymour
Business Manager

QUALITY ASSURANCE
AND EVALUATION

Jerrdean Franklin
Quality Assurance Specialist

Hazel Johnson
Program Evaluation Specialist

Youths, Adolescents and
Young Adults Center

Anthony Fox
Capacity-Building Assistance
Coordinator

Vaughn Smith
Prevention Specialist

Jessica Townsend
Prevention Specialist

YAYA ADVISORY BOARD

William J. Campbell
Houston, Texas

Dr. Mark Colomb
Jackson, Mississippi

Christopher Lane
Washington, D.C.

Erise Williams, Jr.
St. Louis, Missouri

Robert Williams
Oakland, California

S. Alex Williams
Newark, New Jersey

Akoma

Executive Editor: Velesha P. Williams

Editor: Pamela McCoy

Managing Editor: Anthony Fox

Graphic Artist: Lee Campbell II

Akoma is a bi-annual publication of the Jackson State University Mississippi Urban Research Center's Youths, Adolescents and Young Adults Center of the African American Prevention Intervention Network (YAYA/APIN).

Akoma is published as a resource for community-based organizations funded by the Centers for Disease Control and Prevention. Inquiries about articles appearing in this issue or requests for additional information concerning capacity-building assistance should be directed to: Youths, Adolescents and Young Adults Center, Jackson State University, JSU Box 17309, Jackson, MS 39217; Toll-free: 1-866-JSU-MURC; Telephone: (601) 979-1400; Fax: (601) 368-2021.

HIV/AIDS among Youth continued...

the estimated total of 42,514 AIDS diagnoses), and 232 young people with AIDS died [1].

- An estimated 7,761 young people were living with AIDS, a 42% increase since 2000, when 5,457 young people were living with AIDS [1].
- Young people for whom AIDS was diagnosed during 1996–2004 lived longer than persons with AIDS in any other age group except those younger than 13 years. Nine years after receiving a diagnosis of AIDS, 76% of those aged 13–24 were alive, compared with

- 81% of those younger than age 13
- 74% of those aged 25–34
- 70% of those aged 35–44
- 63% of those aged 45–54
- 53% of those aged 55 and older [1].

Since the beginning of the epidemic, an estimated 40,059 young people in the United States had received a diagnosis of AIDS, and an estimated 10,129 young people with AIDS had died. They accounted for about 4% of the estimated total of 944,306 AIDS diagnoses and 2% of the 529,113 deaths of people with AIDS [1].

RISK FACTORS AND BARRIERS
TO PREVENTION

SEXUAL RISK FACTORS

Early age at sexual initiation. According to CDC's Youth Risk Behavioral Survey (YRBS), many young people begin having sexual intercourse at early ages: 47% of high school students have had sexual intercourse, and 7.4% of them reported first sexual intercourse before age 13 [4]. HIV/AIDS education needs to take place at correspondingly young ages, before young people engage in sexual behaviors that put them at risk for HIV infection.

Heterosexual transmission. Young women, especially those of minority races or ethnicities, are increasingly at risk for HIV infection through heterosexual contact. According to data from a CDC study of HIV prevalence among disadvantaged youth during the early to mid-1990s, the rate of HIV prevalence among young women aged

16–21 was 50% higher than the rate among young men in that age group [5]. African American women in this study were 7 times as likely as white women and 8 times as likely as Hispanic women to be HIV-positive. Young women are at risk for sexually transmitted HIV for several reasons, including biologic vulnerability, lack of recognition of their partners' risk factors, inequality in relationships, and having sex with older men who are more likely to be infected with HIV.

MSM. Young MSM are at high risk for HIV infection, but their risk factors and the prevention barriers they face differ from those of persons who become infected through heterosexual contact. According to a CDC study of 5,589 MSM, 55% of young men (aged 15–22) did not let other people know they were sexually attracted to men [6]. MSM who do not disclose their sexual orientation are less likely to seek HIV testing, so if they become infected, they are less likely to know it. Further, because MSM who do not disclose their sexual orientation are likely to have 1 or more female sex partners, MSM who become infected may transmit the virus to women as well as to men. In a small study of African American MSM college students and nonstudents in North Carolina, the participants had sexual risk factors for HIV infection, and 20% had a female sex partner during the preceding 12 months [7].

Sexually transmitted diseases (STDs).

The presence of an STD greatly increases a person's likelihood of acquiring or transmitting HIV [8]. Some of the highest STD rates in the country are those among young people, especially young people of minority races and ethnicities [9].

Substance Abuse

Young people in the United States use alcohol, tobacco, and other drugs at high rates [10]. Both casual and chronic substance users are more likely to engage in high-risk behaviors, such as unprotected sex, when they are under the influence of drugs or alcohol [11]. Runaways and other homeless young people are at high risk for HIV

if they are exchanging sex for drugs or money.

Lack of Awareness

Research has shown that a large proportion of young people are not concerned about becoming infected with HIV [12]. Adolescents need accurate, age-appropriate information about HIV infection and AIDS, including how to talk with their parents or other trusted adults about HIV and AIDS, how to reduce or eliminate risk factors, how to talk with a potential partner about risk factors, where to get tested for HIV, how to use a condom correctly. Information should also include the concept that abstinence is the only 100% effective way to avoid infection.

Poverty and Out-of-School Youth

Nearly 1 in 4 African Americans and 1 in 5 Hispanics live in poverty [13]. The socioeconomic problems associated with poverty, including lack of access to high-quality health care, can directly or indirectly increase the risk for HIV infection [14]. Young people who have dropped out of school are more likely to become sexually active at younger ages and to fail to use contraception [15].

The Coming of Age of HIV-Positive Children

Many young people who contracted HIV through perinatal transmission are facing decisions about becoming sexually active. They will require ongoing counseling and prevention education to ensure that they do not transmit HIV.

PREVENTION

In the United States, the annual number of new HIV infections has declined from a peak of more than 150,000 in the mid-1980s and has stabilized since the late 1990s at approximately 40,000. Populations of minority races or ethnicities are disproportionately affected by the HIV epidemic. To reduce further the

incidence of HIV, CDC announced a new initiative, Advancing HIV Prevention, in 2003. This initiative comprises 4 strategies: making HIV testing a routine part of medical care, implementing new models for diagnosing HIV infections outside medical settings, preventing new infections by working with HIV-infected persons and their partners, and further decreasing perinatal HIV transmission.

Through the Minority AIDS Initiative, CDC explores ways to reduce health disparities in communities made up of persons of minority races or ethnicities who are at high risk for HIV. These funds are used to address the high-priority HIV prevention needs in such communities.

CDC provides 9 awards to community-based organizations (CBOs) that focus primarily on youth and provides indirect funding through state, territorial, and local health departments to organizations serving youth. Of these 9 awards, 5 are focused on African Americans, 3 on Hispanics, 1 on Asians and Pacific Islanders, and 1 on whites. The following are some CDC-tested provide for youth.

- Teens Linked to Care is focused on young people aged 13–29 who are living with HIV.
- Street Smart is an HIV/AIDS and STD prevention program for runaway and homeless youth.
- PROMISE (Peers Reaching Out and Modeling Intervention Strategies for HIV/AIDS Risk Reduction in their Community) is a community-level HIV prevention intervention that relies on role-model stories and peers from the community.
- Adult Identity Mentoring project, which encourages students to articulate personal goals and then teaches them the skills required to achieve those goals, can be effective in helping at-risk youth

CDC research has shown that early, clear parent-child communication regarding values and expectations about sex is an important step in helping adolescents delay sexual initiation and make responsible decisions about sexual behaviors later in life. Parents are in a unique position to engage their children in conversations about HIV, STD, and teen pregnancy prevention because the conversations can be ongoing and timely [17].

Schools also can be important partners for reaching youth before high-risk behaviors are established, as evidenced by the YRBS finding that 88% of high school students in the United States reported having been taught about AIDS or HIV infection in school.

Source: <http://www.cdc/hiv/resources/factsheets/youth.htm>

REFERENCES

1. CDC *HIV/AIDS Surveillance Report*, 2004. Vol. 16. Atlanta: US Department of Health and Human Services, CDC; 2005:1–46.
2. CDC. *HIV Prevention in the Third Decade*. Atlanta: US Department of Health and Human Services, CDC; 2005.
3. CDC. HIV incidence among young men who have sex with men—seven US cities, 1994–2000. *MMWR* 2001;50:440–444.
4. CDC. Youth Risk Behavior Surveillance—United States, 2003. *MMWR* 2004;53(SS-2):1–29.
5. Valleroy LA, MacKellar DA, Karon JM, Janssen RS, Hayman DR. HIV infection in disadvantaged out-of-school youth: prevalence for U.S. Job Corps entrants, 1990 through 1996. *Journal of Acquired Immune Deficiency Syndromes* 1998;19:67–73.
6. CDC. HIV/STD risks in young men who have sex with men who do not disclose their sexual orientation—six US cities, 1994–2000. *MMWR* 2003;52:81–85.
7. CDC. HIV transmission among black college student and non-student men who have sex with men—North Carolina, 2003. *MMWR* 2004;53:731–734.
8. Fleming DT, Wasserheit JN. From epidemiological synergy to public health policy and practice: the contribution of other sexually transmitted diseases to sexual transmission of HIV infection. *Sexually Transmitted Infections* 1999;75:3–17. 9.
9. CDC. *Sexually Transmitted Disease Surveillance*.